

# Morphometrics, length-weight relationship and condition of *Rastrineobola argentea* (Pellegrin 1904) in the Winam Gulf of Lake Victoria (Kenya)

J. O. MANYALA,<sup>1</sup> E. VANDEN BERGHE<sup>2</sup> AND S. DADZIE<sup>3</sup>

1. Moi University, Fisheries Department, P. O. Box 3900, Eldoret, Kenya.

2. Regional Co-operation in Scientific Exchange in the Western Indian Ocean Region (RECOSCIX-WIO), P. O. Box 95832, Mombasa, Kenya.

3. Department of Zoology, Kuwait University, P. O. Box 5969, Safat 13069, Kuwait.

## ABSTRACT

Correlation between total length (TL), fork length (FL) and standard length (SL) of *Rastrineobola argentea* (Pellegrin 1904) in the Winam Gulf of Lake Victoria indicate that  $FL = 0.92 TL - 0.74$  and  $SL = 0.90 TL - 1.74$ . Length-weight relationship of log-transformed data shows that the slopes of the regression lines were 3.06 to 3.22 for juveniles, 2.70 to 3.05 for males and 3.24 to 3.71 for females. These slopes were significantly different between groups at  $\alpha = 0.05$ . The Fulton's condition factor (K) was highest in December (1.019-1.073) and March/April (1.015-1.030) but lowest in June (1.00-1.025) for all stations. Significant differences between groups demands for the use of different growth models for juveniles, males and females especially for the von Bertalanffy growth equation which uses length-weight relationship. Observed cyclic variations in condition factor suggests two peak breeding seasons for this species in the Winam Gulf. The practical implication of these results in stock assessment using length-based fish stock assessment methods is briefly discussed.

## INTRODUCTION

The family Cyprinidae are known as typical marine family. However, several members of the family are lake inhabitants. Some of these lacustrine cyprinids have not only adopted a pelagic way of life, but also their shape and silvery camouflage (GREENWOOD, 1966; WANINK, 1989) resemble marine pelagic fish such as herring (*Clupea sp.*). The pelagic cyprinids endemic to some East and Central African lakes were previously placed in one genus, *Engraulicypris*, until a revision by HOWES (1980). In Lakes Victoria, Kyoga and Nabugabo, this group of fishes is represented by *Rastrineobola argentea* (Pellegrin) according to the current classification by Howes (*op. cit.*) and *Chelaethiops sp.* in Lake Tanganyika which is of no economic value (OKEDI, 1981B; WANINK, 1989). *R. argentea* is locally known as "omena" in Kenya, "mukene" in Uganda and "nsalali" in Tanzania (GRAHAM, 1929; GREENWOOD, 1966). The name "dagaa" became popular in the early 1970s when a light fishery for this species reached commercial significance. The fishermen of Lake Victoria adopted this name from the clupeids of Lake Tanganyika which were also caught by light attraction (OKEDI, 1981A; 1981B; WANINK, 1989).

Fish stock assessment has been carried out in the Winam Gulf of Lake Victoria using bottom trawls (KUDHONGANIA and CORDONE, 1974; MARTEN *et al.*, 1976; BENDA, 1981; MULLER and BENDA, 1981), a catch assessment survey on the artisanal fishery (RABUOR, 1988) and length-frequency analysis (GETABU, 1988; ASILA and OGARI, 1988). Very little scientific information on the biology and ecology of *Rastrineobola argentea* is available. The earlier works of GRAHAM (1929); GREENWOOD (1966) were brief summaries covering all the known fish species of Lake Victoria at that time. The preliminary observations by OKEDI (1973) on the breeding ecology and fecundity of *Rastrineobola* in Lake Victoria may be considered as the pioneer work on the biology and ecology of the species in Lake Victoria. Recent work on *R. argentea* includes those of WANINK (1989); MANYALA (1991); CHITAMWEBWA (1992); WANDERA (1992); KATUNZI (1992); MANYALA *et al.* (1992) on the general biology and ecology of the species in Lake Victoria. MANNINI (1992) reviewed the characteristics of *R. argentea* in Lake Victoria and compared it to other small pelagics from other African Great Lakes.



The correlation between the different forms of length measurements are shown in Fig. 2 while the details of these regression analyses are given in Table 1. The correlation coefficients in all the

analyses were very high ( $r^2 = 0.90 - 0.99$ ) while the standard error of each coefficient was very low (0.01).




Fig. 2. Correlation between total length (TL), fork length (FL) standard length (SL) of *R argentea* in the Winam Gulf.

Table 1: Correlation between total length (TL), fork length (FL) and standard length (SL) of *R. argentea* in the Winam Gulf of Lake Victoria.

Station	Indep. variable	Dependent variable	Coeff. $\pm$ SE	Const	$r^2$	n
Sango	TL	FL	$0.9 \pm 0$	-0.2	0.90	249
	TL	SL	$0.8 \pm 0$	-0.3	0.90	249
	FL	SL	$0.9 \pm 0$	-0.4	0.90	249
Rakwaro	TL	FL	$0.9 \pm 0,01$	-0.8	0.96	285
	TL	SL	$0.9 \pm 0.01$	-0.8	0.96	285
	FL	SL	$0.9 \pm 0.01$	-0.5	0.96	285
Obaria	TL	FL	$0.9 \pm 0$	-1.0	0.99	238
	TL	SL	$0.9 \pm 0$	-4.0	0.99	238
	FL	SL	$0.9 \pm 0$	-2.0	0.99	238
Ngegu	TL	FL	$0.9 \pm 0.01$	-1.3	0.98	239
	TL	SL	$0.9 \pm 0.01$	-1.3	0.98	239
	FL	SL	$0.9 \pm 0.01$	-0.2	0.99	239
Luanda	TL	FL	$0.9 \pm 0.01$	-0.6	0.98	220
	TL	SL	$0.9 \pm 0.01$	-2.4	0.98	220
	FL	SL	$0.9 \pm 0.01$	-2.7	0.98	220

**Table 2.** Correlation between Log TL and Log W of *R. argentea* in the Winam Gulf. Similar superscripts indicate no significant differences

Station	Length-weight relationship	n	r <sup>2</sup>
<b>Juveniles:</b>			
Sango <sup>1</sup>	Log W = 3.06 Log TL - 5.42	732	0.60
Rakwaro <sup>1</sup>	Log W = 3.12 Log TL - 5.60	708	0.68
Obaria <sup>1</sup>	Log W = 3.10 Log TL - 5.50	663	0.72
Ngegu <sup>1</sup>	Log W = 3.20 Log TL - 5.84	479	0.66
Luanda <sup>1</sup>	Log W = 3.22 Log TL - 6.00	456	0.84
<b>Males:</b>			
Sango <sup>2</sup>	Log W = 2.70 Log TL - 4.69	754	0.60
Rakwaro <sup>2</sup>	Log W = 2.81 Log TL - 4.70	977	0.69
Obaria <sup>2</sup>	Log W = 3.05 Log TL - 4.90	815	0.64
Ngegu <sup>2</sup>	Log W = 2.94 Log TL - 5.00	822	0.67
Luanda <sup>2</sup>	Log W = 2.75 Log TL - 4.76	985	0.70
<b>Females:</b>			
Sango <sup>3</sup>	Log W = 3.60 Log TL - 6.20	1266	0.66
Rakwaro <sup>3</sup>	Log W = 3.65 Log TL - 6.41	1348	0.72
Obaria <sup>3</sup>	Log W = 3.71 Log TL - 6.50	1144	0.68
Ngegu <sup>3</sup>	Log W = 3.40 Log TL - 5.60	1219	0.68
Luanda <sup>3</sup>	Log W = 3.24 Log TL - 5.45	1337	0.61
F(14,13675)	6.324*		

**Length-weight relationship**

The mean length-weight relationship of the species from each of the five stations is presented in Table 2. The graphical representation of length-weight relationship is shown in Fig. 3 before any transformation and Fig. 4 after log-log transformation. An analysis of covariance (ANCOVA) revealed significant differences between the

slopes of the regression lines (Table 3) while multiple range test, Student Newman-Keuls (SNK) test indicated that there were significant differences between the slopes for juveniles, males and females but no differences within sexes from different sampling stations. These results suggest the use of different growth models for the three groups considered in this study.



**Fig. 3.** An example of length-weight relationship of *R. argentea* in the Winam Gulf from Sango.**Fig. 4.** Correlation between Log L and Log W of *R. argentea* in the Winam Gulf for samples taken from Sango.

Table 3. ANOVA of Fultons condition factor of *R. argentea* in the Winam Gulf of Lake Victoria

Month	Sango	Rakwaro	Obaria	Ngegu	Luanda
August	1.015	1.011	1.011	1.024	1.015
December	1.073	1.048	1.020	1.024	1.019
January	1.048	1.011	1.019	1.017	1.017
February	1.021	1.013	1.011	1.019	1.016
March	1.036	1.035	1.045	1.025	1.028
April	1.015	1.023	1.024	1.030	1.029
May	1.015	1.023	1.024	1.025	1.027
June	1.000	1.023	1.002	1.025	1.022
July	1.013	1.028	1.013	1.022	1.020
F(4,32)	0.113 ns				stations
F(8,32)	4.323 *				months

### Fultons condition factor (K)

There was an initial gain in condition, reaching a peak in December followed by a decline up to February. This decline was followed by a second gain in condition reaching a peak in March/April. Thereafter, there was a general decline in condition up to July. Maximum K value recorded in December was 1.074 while that of March was 1.052. The lowest condition recorded in June was 1.000 and July was 1.006. All stations showed a similar pattern of relative condition factor by months as shown in Fig. 5. Analysis of variance indicated real differences in the Fulton's condition factor (K) in different months but there were no significant differences between stations (Table 4). Weight measurements could not be taken from September to December.

### DISCUSSION AND CONCLUSION

The linear equations for the interconversion of different forms of length measurements determined by this study are close to those of WANINK (1989) in the Mwanza Gulf of Lake Victoria and had very high correlation coefficients. The sample size for the Mwanza Gulf was relatively small (45-52) while the large sample size for this study (220-285) confirms these previous results. These values can thus be used to convert one form of length measurement to the other for comparison due to individual preferences in the form of length measurements.




Fig. 5. Fulton's condition factor (K) of *R. argentea*. Vertical bars shows 2SD.

Although no values are available for direct comparison, the length-weight relationships derived by this study can form the basis of future work on this species. Information on length-weight relationship serves two purposes: a) the first is a mathematical description of the relationship between length and weight so that one of the parameters can be converted into the other. b) the second measures the variation from expected weight-for-length of the individual fish which is used in the calculations of condition factor (DADZIE *et al.*, 1979; DADZIE, 1985). For the length-based fish stock assessment methods, it is important to consider the length-weight relationship in totality before assuming isometric growth. The exponent in the length-weight relationship seemed to be quite high in this study although the growth of *R. argentea* was assumed to be isometric by WANINK (1989). Such a parallel relationship could be attributed to sampling errors in this study. Since weight of the specimens were not recorded from September to November, the observed changes only indicate a general trend in the Fulton's condition factor. The observed changes in Fulton's condition factor could however be attributed to development of the gonadal material during or just prior to the peak breeding seasons. Maximum condition factor incidentally fell within the established rainy seasons for the region and other fish species in Lake Victoria have been established to breed during the same time of the year (OKEDI, 1969; MARTEN, 1979; MULLER *et al.*, 1982; RINNE and WANJALA, 1983; OCHUMBA and MANYALA, 1992).

#### ACKNOWLEDGEMENT

We wish to thank the Director, KMFRI, Dr. E. Okemwa and the Deputy Director (Inland Waters), KMFRI, Mr. J. Ogari for supporting this work and availing both laboratory space and field facilities for the study. We extend our sincere thanks to the Kenyan Government for awarding Mr. J. Manyala a scholarship which enabled us complete this work as a part of an M. Sc. Thesis. We would also like to thank the teaching staff at the University of Nairobi (Zoology Department) who gave valuable advice on this study and many other people who helped in their own way to see this work to completion. We would also like to acknowledge financial and technical assistance given by the International Foundation for Science

(IFS) to continue with research in this area and process this manuscript.

#### REFERENCES

- Asila, A. A. and J. Ogari (1988). Growth parameters and mortality of Nile perch (*Lates niloticus*) estimated from length-frequency in the Nyanza Gulf (Lake Victoria). *FAO Fish. Rep.* 389:272-287.
- Bagenal, T. B. and E. Braum (1971). Egg and early life history. *In*. Methods of assessment of fish production in fresh waters. E. W. Ricker (ed.). 2nd. edition, IBP handbook No. 3.
- Benda, R. S. (1981). A comparison of bottom trawl catch rates in the Kenyan waters of Lake Victoria. *J. Fish Biol.* 18:13-24.
- Chitamwebwa, D. B. R. (1988). Historical development of dagaa fishery policy and planning. Seminar paper, University of Dar-es-Salaam, 2-4 May 1988. 11pp.
- Copley, H. (1953). The tilapia fishery of the Kavirondo Gulf. *J. E.. African Nat. Hist. Soc.* 94: 1-5.
- Dadzie, S. (1985). Seasonal changes in length-weight relationship, reproduction and condition of the African carp, *Labeo cylindricus* (Peters) in a tropical man-made lake, Lake Kamburu, Kenya. *Kenya Journal of Science and Technology, Series B* (1985) (2): 79-88.
- Dadzie, S., R. S. Odingo and A. Ongoma (1979). Aquatic ecology and fisheries. *In*. An African Dam: Ecological surveys of the Kamburu/Gitaru hydro-electric dam area, Kenya. Edited by R. S. Odingo. *Ecol. Bull. (Stockholm)* 29: 65-84.
- Getabu, A. (1988). Aspects of the Lake Victoria fisheries, with emphasis on *Oreochromis niloticus* and *Alestes sadleri* from the Nyanza Gulf. *FAO Fish. Rep.* 389: 416-431.
- Graham, M. (1929). The Victoria Nyanza and its fisheries. Crown agents, London.
- Greenwood, P. H. (1966). The fishes of Uganda. 2nd. Ed. The Uganda Society, Kampala.
- Hopson, J. (1972). Breeding and growth in two populations of *Alestes baremose* (Joannis.) (Pisces: Characidae) from the northern basin of Lake Chad. Overseas research publication No. 22: 50 pp. HMSO London.
- Hopson, J. (1975). Preliminary observations on the biology of *Alestes baremose* (Joannis.) in Lake Rudolf. *Symp. Hydrobiol. Fish. of L. Rudolf, Molo, 25-29 May 1975.*
- Howes, G. (1980). The anatomy, phylogeny and classification of barilaine cyprinid fishes. *Bull. Brit. Mus. Nat. Hist. (Zool.)* 37: 129-198.

- Howes, G. (1980). The anatomy, phylogeny and classification of bariliine cyprinid fishes. *Bull. Brit. Mus. Nat. Hist. (Zool.)* 37: 129-198.
- Katunzi, E. F. B. (1992). Biological and fishery aspects of *Rastrineobola argentea* in the southern part of Lake Victoria. In: Mannini, P. (ed.), The Lake Victoria Dagaa (*Rastrineobola argentea*). Report of the First Meeting of the Working Group on Lake Victoria, 9-11 December 1991, Kisumu, Kenya. *UNDP/FAO Regional Project for Inland Fisheries Planning (IFIP) RAF/87/099-TD/92 (En.)*: 51-57.
- Kudhongania, A. W. and A. J. Cordone (1974). Bathospacial distribution pattern and biomass estimates of major demersal fishes in Lake Victoria. *Afr. J. Trop. Hydrobiol. Fish.* 3: 15-31.
- Mannini, P. (1992). Some characteristics of small pelagic fish species and possible affinities with the population of Lake Victoria Dagaa (*Rastrineobola argentea*). In: Mannini, P. (ed.), The Lake Victoria Dagaa (*Rastrineobola argentea*). Report of the First Meeting of the Working Group on Lake Victoria, 9-11 December 1991, Kisumu, Kenya. *UNDP/FAO Regional Project for Inland Fisheries Planning (IFIP) RAF/87/099-TD/92 (En.)*: 62-78.
- Manyala, J. O. (1991). Population dynamics of *Rastrineobola argentea* (Pellegrin 1904) (Pisces: Cyprinidae) in the Winam Gulf of Lake Victoria (Kenya). *M. Sc. Thesis, University of Nairobi*, 107 pp.
- Manyala, J. O. (1992). Growth, mortality and gear selection of *Rastrineobola argentea* (Pellegrin) in the Winam Gulf of Lake Victoria, Kenya. p 246-256. In: B. E. Marshall and R. Mubamba (eds.). Symposium on the biology, stock assessment and exploitation of small pelagic fish species in the African Great Lakes region. 24th. - 28th. November 1992. Bujumbura, Burundi. *CIFA Occasional Paper No. 19*.
- Manyala, J. O., C. O. Nyawade and C. O. Rabuor (1992). The Dagaa (*Rastrineobola argentea* Pellegrin) fishery in the Kenyan waters of Lake Victoria: A national review and proposal for future research. In: Mannini, P. (ed.), The Lake Victoria Dagaa (*Rastrineobola argentea*). Report of the First Meeting of the Working Group on Lake Victoria, 9-11 December 1991, Kisumu, Kenya. *UNDP/FAO Regional Project for Inland Fisheries Planning (IFIP) RAF/87/099-TD/92 (En.)*: 18-35.
- Marten, G. G. (1979). Impact of fishing in the inshore fishery of Lake Victoria (East Africa). *Journal of the Fisheries Research Board of Canada* 36: 891-900.
- Marten, G. G., B. Wanjala and L. T. Galuka (1976). Exploratory trawling of Lake Victoria fishery in Kenya during 1975. *EAFFRO manuscript*, 19 pp.
- Muller, R. G. and R. S. Benda (1981). A comparison of bottom stock densities in the inner Kavirondo Gulf of Lake Victoria. *J. Fish Biol.* 19: 339-401.
- Muller, R. G., R. S. Benda and T. G. Powell (1982). Exploitation of fish in the inner Kavirondo Gulf of Lake Victoria, Kenya based on tag returns. *Journal of Fish Biology* 20: 667-672.
- Ochumba, P. B. O. and J. O. Manyala (1992). Distribution of fishes along the Sondu-Miriu River of Lake Victoria, Kenya with special reference to upstream migration, biology and yield. *Journal of Aquaculture and Fisheries Management* 23: 701-719.
- Okedi, J. (1973). Preliminary observations on *Engraulicypris argentea* from Lake Victoria. *EAFFRO annual report (1973)*.
- Okedi, J. (1981a). Integrated management of the "dagaa" fishery of Lake Victoria. In: Proceedings of the workshop of the Kenya Marine and Fisheries Research Institute on aquatic resources of Kenya. 13-19 July 1981.
- Okedi, J. (1981b). The *Engraulicypris* "dagaa" fishery of Lake Victoria: with special reference to the southern waters of the Lake. In: Proceedings of the workshop of the Kenya Marine and Fisheries Research Institute on aquatic resources of Kenya. 13-19 July 1981.
- Rabuor, C. O. (1988). First report of the fish stock assessment on artisanal fishery of the Winam Gulf of Lake Victoria (Kenya). *K.M.F.R.I. Tech. Rep. 20 February 1988*.
- Rinne, J. N. and B. Wanjala (1983). Maturity, fecundity and breeding seasons of major catfishes (Suborder: Siluroidae) in Lake Victoria, East Africa. *Journal of Fish Biology* 23: 357-363.
- Wandera, S. B. (1990). The population exploitation of small fishes of the Great Lakes with reference to the mukene (*Rastrineobola argentea*) fishery of northern waters of Lake Victoria. In: Fisheries of African Great Lakes. Research papers presented at the International Symposium on Resource Use and Conservation of African Great Lakes. Bujumbura, 29th. November - 2nd. December 1989. *International Agricultural Centre, Wageningen, The Netherlands, Fisheries and Aquaculture Unit. Occasional Paper 3*: 67-74.
- WANDERA, S. B. (1992). A study of *Rastrineobola argentea* in the Uganda lakes. In: Mannini, P. (ed.), The Lake Victoria Dagaa (*Rastrineobola*

- Wandera, S. B. (1992). A study of *Rastrineobola argentea* in the Uganda lakes. In: Mannini, P. (ed.), The Lake Victoria Dagaa (*Rastrineobola argentea*). Report of the First Meeting of the Working Group on Lake Victoria, 9-11 December 1991, Kisumu, Kenya. *UNDP/FAO Regional Project for Inland Fisheries Planning (IFIP) RAF/87/099-TD/92 (En.)*: 36-50.
- Wanink, J. H. (1989). The ecology of dagaa, *Rastrineobola argentea* (Pellegrin) 1904. Report of the Haplochromis Ecology Survey Team (HEST) operating in Lake Victoria. HEST/TAFIRI/ FAO/DANIDA workshop on the fish stock in Lake Victoria. January/February 1989, Mwanza, Tanzania.
- Zar, J. H. (1984). Biostatistical analysis. Prentice-Hall International, Inc., Englewood Cliffs, New Jersey. 718 pp.